Asarina

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.23**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values
Temperature indicator value: **8**
Disturbance indicator values
Disturbance frequency: **0.36**
Disturbance frequency (herb layer): **0.67**
Disturbance severity: **0.26**
Disturbance severity (herb layer): **0.12**
Mowing frequency: **0.04**
Grazing pressure: **0.1**
Soil disturbance: **0.06**

Habitat and sociology

Syntaxon
Diagnostic species of phytosociological classes: **KC (ASP) *Asplenietea trichomanis***
EUNIS habitat
Diagnostic species of EUNIS habitats: **U34 Mediterranean siliceous inland cliff**
Broad habitat
Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution

Continentality: **2**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz