Convallaria

Habitus and growth type

Plant height [m]: **0.2**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **28.97**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **235.9**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **4.9**
Reaction indicator value: **5.6**
Nutrient indicator value: **3.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.05**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.13**
Mowing frequency: **0.02**
Grazing pressure: **0.2**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***, **HA (POP) *Alno glutinosae-Populetea albae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T1E Carpinus and Quercus mesic deciduous forest**

Constant species of EUNIS habitats: **T1E Carpinus and Quercus mesic deciduous forest**, **T35 Temperate continental Pinus sylvestris forest**, **T3F Dark taiga**, **T3G Pinus sylvestris light taiga**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz