

Limbarda

Habitus and growth type

Plant height [m]: **0.52**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **June-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.34**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.9**

Reaction indicator value: **8.3**

Nutrient indicator value: **6.2**

Salinity indicator value: **6**

Disturbance indicator values

Disturbance frequency: **1.27**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.04**

Grazing pressure: **0.19**

Soil disturbance: **0.28**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed](#), [MA253 Mediterranean mid-low saltmarsh](#)

Constant species of EUNIS habitats: [MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed](#), [MA253 Mediterranean mid-low saltmarsh](#), [N31 Atlantic and Baltic rocky sea cliff and shore](#), [N32 Mediterranean and Black Sea rocky sea cliff and shore](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Coastal cliff, Saline vegetation**