Parasenecio

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **53.19**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Disturbance indicator values

Disturbance frequency: **0.04**
Disturbance frequency (herb layer): **1.32**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.02**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BB (ASA) *Asaro europaei-Abietetea sibiricae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3H Larix light taiga**

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz