Sibbaldianthe

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.5**
Moisture indicator value: **4.3**

Disturbance indicator values

Disturbance frequency: **1.41**
Disturbance frequency (herb layer): **1.82**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.66**
Mowing frequency: **0.67**
Grazing pressure: **0.18**
Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **DB (FEP) *Festuco-Puccinellietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S68 Semi-desert sand dune with sparse scrub**
Constant species of EUNIS habitats: **S68 Semi-desert sand dune with sparse scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Saline vegetation, Semi-desert**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz