

# ***Stratiotes***

## **Habitus and growth type**

Plant height [m]: **0.23**

Life span: **Perennial**

Life form: **Hydrophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **23.51**

## **Flower**

Flowering period: **June-July**

## **Fruit, seed and dispersal**

Dispersal mode: **Hydrochory**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **6.2**

Moisture indicator value: **11.1**

Reaction indicator value: **7.3**

Nutrient indicator value: **6.2**

Salinity indicator value: **0.2**

### Disturbance indicator values

Disturbance frequency: **1.51**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.02**

Grazing pressure: **0.01**

Soil disturbance: **0.01**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [NA \(LEM\) Lemnetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [P3c Fresh-water large pleustophyte vegetation](#)

Constant species of EUNIS habitats: [P3c Fresh-water large pleustophyte vegetation](#)

Dominant species of EUNIS habitats: [P3c Fresh-water large pleustophyte vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic**

## Distribution

Continentality: **6**

Continentality amplitude: **6**