

# *Cyanus triumfettii* aggr.

## Habitus and growth type

Plant height [m]: **0.31**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **2.39**

## Fruit, seed and dispersal

Seed mass [mg]: **9.82**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.5**

Moisture indicator value: **3.8**

Reaction indicator value: **7.2**

Nutrient indicator value: **3.1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.57**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.18**

Grazing pressure: **0.27**

Soil disturbance: **0.13**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*S24 Subalpine genistoid scrub of the Amphi-Adriatic region\*\*](#)

Constant species of EUNIS habitats: [\*\*S24 Subalpine genistoid scrub of the Amphi-Adriatic region\*\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**