# *Arenaria ciliata* aggr.

### Habitus and growth type

Plant height [m]: **0.06**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

### Flower

Flowering period: **June-August**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **8.8**
Temperature indicator value: **1.6**
Moisture indicator value: **4.8**
Reaction indicator value: **7.5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **1.47**
Disturbance frequency (herb layer): **1.64**
Disturbance severity: **0.29**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.03**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Grazing pressure: **0.26**
Soil disturbance: **0.12**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R44 Arctic-alpine calcareous grassland**, **S21 Subarctic and alpine dwarf Salix scrub**

### Broad habitat

Occurrence in broad habitats: **Spring, Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **6**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*