

Crepis foetida aggr.

Habitus and growth type

Plant height [m]: **0.23**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **16.08**

Flower

Flowering period: **May-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.61**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**

Moisture indicator value: **3.6**

Reaction indicator value: **6.6**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.27**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.62**

Mowing frequency: **0.55**

Grazing pressure: **0.27**

Soil disturbance: **0.36**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [**S72 Eastern Mediterranean spiny heath \(phrygana\)**](#)

Constant species of EUNIS habitats: [**S72 Eastern Mediterranean spiny heath \(phrygana\)**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**