

# *Oenothera biennis* aggr.

## Habitus and growth type

Plant height [m]: **1.58**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **15.7**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.52**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **6.7**

Moisture indicator value: **4**

Reaction indicator value: **6.6**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.31**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.6**

Mowing frequency: **0.6**

Grazing pressure: **0.19**

Soil disturbance: **0.34**

## **Habitat and sociology**

### **Syntaxon**

Diagnostic species of phytosociological classes: [QF \(ART\) Artemisietea vulgaris](#)

### **Broad habitat**

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree),**

**Synanthropic**

## **Distribution**

Continentality amplitude: **5**