Poa badensis aggr.

Habitus and growth type

Plant height [m]: **0.23**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.28**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**
Temperature indicator value: **6.9**
Moisture indicator value: **2.4**
Reaction indicator value: **8.3**
Nutrient indicator value: **1.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.4**
Disturbance frequency (herb layer): **1.67**
Disturbance severity: **0.36**
Disturbance severity (herb layer): **0.34**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.14**
Grazing pressure: **0.31**
Soil disturbance: **0.18**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R16 Perennial rocky grassland of Central and South-Eastern Europe**, **R17 Heavy-metal dry grassland of the Balkans**
Constant species of EUNIS habitats: **R16 Perennial rocky grassland of Central and South-Eastern Europe**, **R17 Heavy-metal dry grassland of the Balkans**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz