Poa pratensis aggr.

Habitus and growth type

Plant height [m]: **0.65**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **21.55**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.25**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic, Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**
Temperature indicator value: **5.1**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.12**
Disturbance frequency (herb layer): **1.87**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.48**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.75**
Grazing pressure: **0.28**
Soil disturbance: **0.3**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R21 Mesic permanent pasture of lowlands and mountains**, **R22 Low and medium altitude hay meadow**, **R34 Submediterranean moist meadow**, **S36 Low steppic scrub**
Constant species of EUNIS habitats: **MA222 Atlantic upper saltmarsh**, **MA232 Baltic coastal meadow**, **N18 Atlantic and Baltic coastal Empetrum heath**, **N1A Atlantic and Baltic coastal dune scrub**, **N1D Atlantic and Baltic broad-leaved coastal dune forest**, **N1H Atlantic and Baltic moist and wet dune slack**, **R1A Semi-dry perennial calcareous grassland (meadow steppe)**, **R1B Continental dry grassland (true steppe)**, **R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**, **R21 Mesic permanent pasture of lowlands and mountains**, **R22 Low and medium altitude hay meadow**, **R23 Mountain hay meadow**, **R34 Submediterranean moist meadow**, **R35 Moist or wet mesotrophic to eutrophic hay meadow**, **R51 Thermophilous forest fringe of base-rich soils**, **R54 Pteridium aquilinum vegetation**, **S32 Temperate Rubus scrub**, **S35 Temperate and submediterranean thorn scrub**, **S36 Low steppic scrub**, **V38 Dry perennial anthropogenic herbaceous vegetation**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Coastal cliff, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz