

Poa pratensis aggr.

Habitus and growth type

Plant height [m]: **0.65**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **21.55**

Flower

Flowering period: **April-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.25**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic, Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **5.1**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.12**

Disturbance frequency (herb layer): **1.87**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.75**

Grazing pressure: **0.28**

Soil disturbance: **0.3**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R22 Low and medium altitude hay meadow](#)

Constant species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#), [MA232 Baltic coastal meadow](#), [N15 Atlantic and Baltic coastal dune grassland \(grey dune\)](#), [N18 Atlantic and Baltic coastal Empetrum heath](#), [N19 Atlantic coastal Calluna and Ulex heath](#), [N1A Atlantic and Baltic coastal dune scrub](#), [N1D Atlantic and Baltic broad-leaved coastal dune forest](#), [N1E Black Sea broad-leaved coastal dune forest](#), [N1H Atlantic and Baltic moist and wet dune slack](#), [N35 Mediterranean and Black Sea soft sea cliff](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1B Continental dry grassland \(true steppe\)](#), [R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils](#), [R21 Mesic permanent pasture of lowlands and mountains](#), [R22 Low and medium altitude hay meadow](#), [R23 Mountain hay meadow](#), [R34 Submediterranean moist meadow](#), [R35 Moist or wet mesotrophic to eutrophic hay meadow](#), [R51 Thermophilous forest fringe of base-rich soils](#), [R54 Pteridium aquilinum vegetation](#), [S32 Temperate Rubus scrub](#), [S35 Temperate and submediterranean thorn scrub](#), [S36 Low steppic scrub](#), [V38 Dry perennial anthropogenic herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: Coastal saltmarsh, Coastal beach, dune or shingle, Coastal cliff, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic