Sorbus aria aggr.

Habitus and growth type

Plant height [m]: **9.5**
Life span: **Perennial**
Life form: **Phanerophyte, Tree, Shrub**
Spinescence: **not spinescent**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **16.06**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **334.49**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**
Temperature indicator value: **6**
Moisture indicator value: **3.9**
Reaction indicator value: **7.2**
Nutrient indicator value: **2.9**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.05**
Disturbance frequency (herb layer): **1.27**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.13**
Mowing frequency: **0.01**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**, **T3C Taxus baccata forest**
Constant species of EUNIS habitats: **S24 Subalpine genistoid scrub of the Amphi-Adriatic region**, **S52 Submediterranean pseudomaquis**, **T17 Fagus forest on non-acid soils**, **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1D Southern European mountain Betula and Populus tremula forest on mineral soils**, **T1F Ravine forest**, **T32 Temperate mountain Abies forest**, **T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**, **T37 Mediterranean montane Pinus sylvestris-Pinus nigra forest**, **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**, **T3C Taxus baccata forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **3**
Continentality amplitude: **3**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz