Utricularia vulgaris aggr.

Habitus and growth type

Plant height [m]: **1.3**
Life span: **Perennial**
Life form: **Hydrophyte**

Leaf

Specific leaf area [mm^2/mg]: **65.66**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**
Temperature indicator value: **5.8**
Moisture indicator value: **11.8**
Reaction indicator value: **5.4**
Nutrient indicator value: **3.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.52**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.62**
Disturbance severity: **0.26**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.12**
Grazing pressure: **0.04**
Soil disturbance: **0.06**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **P3c Fresh-water large pleustophyte vegetation, V14 Inundated or inundatable cropland, including rice fields**
Constant species of EUNIS habitats: **P3c Fresh-water large pleustophyte vegetation, V14 Inundated or inundatable cropland, including rice fields**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz