

Veronica hederifolia aggr.

Habitus and growth type

Plant height [m]: **0.09**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **28.82**

Flower

Flowering period: **January-May, December**

Fruit, seed and dispersal

Seed mass [mg]: **3.48**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **4.5**

Reaction indicator value: **6.6**

Nutrient indicator value: **6.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.54**

Disturbance frequency (herb layer): **1.64**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.61**

Mowing frequency: **0.27**

Grazing pressure: **0.16**

Soil disturbance: **0.54**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#), [V15 Bare tilled, fallow or recently abandoned arable land](#)

Constant species of EUNIS habitats: [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#), [V15 Bare tilled, fallow or recently abandoned arable land](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**