Vicia cracca aggr.

Habitus and growth type

Plant height [m]: **0.78**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **26.47**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **20.59**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry, Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral, Alkaline**
Nutrient relationship: **Oligotrophic, Mesotrophic**
Salinity relationship: **Non-saline, Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **5**
Moisture indicator value: **4**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **1**

Habitat and sociology

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz