

# ***Taraxacum sect. Alpina***

## **Habitus and growth type**

Plant height [m]: **0.13**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **27.4**

## **Flower**

Flowering period: **May-August**

## **Fruit, seed and dispersal**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic, Wet**

Substrate reaction relationship: **Acidic, Alkaline**

Nutrient relationship: **Dystrophic, Oligotrophic, Mesotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8**

Temperature indicator value: **2.1**

Moisture indicator value: **6.3**

Reaction indicator value: **6.5**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.59**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.06**

Grazing pressure: **0.22**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) \*Thlaspietea rotundifolii\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R41 Snow-bed vegetation](#)

Constant species of EUNIS habitats: [R41 Snow-bed vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Mire, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **3**

Continentality amplitude: **3**