

# *Acer platanoides*

## Habitus and growth type

Plant height [m]: **22.87**

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **19.66**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **122.63**

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **4, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.7**

Temperature indicator value: **5.8**

Moisture indicator value: **5.6**

Reaction indicator value: **6.5**

Nutrient indicator value: **6.3**

Salinity indicator value: **0**

## Disturbance indicator values

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.15**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) Carpino-Fagetea sylvaticae](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T1F Ravine forest](#)

Constant species of EUNIS habitats: [T17 Fagus forest on non-acid soils](#), [T1E Carpinus and Quercus mesic deciduous forest](#), [T1F Ravine forest](#)

### Broad habitat

Occurrence in broad habitats: **Forest**

## Distribution

Continentality: **5**

Continentality amplitude: **4**