Acer sempervirens

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Fruit, seed and dispersal

Seed mass [mg]: **26.74**
Dispersal mode: **Anemochory**
Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5**
Moisture indicator value: **5**
Reaction indicator value: **7**
Nutrient indicator value: **4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.1**
Disturbance frequency (herb layer): **1.46**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0.03**
Grazing pressure: **0.3**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3D Mediterranean Cupressaceae forest**

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz