Agrimonia procera

Habitus and growth type

Plant height [m]: **0.88**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **19.52**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **26.39**
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.6**
Moisture indicator value: **5.3**
Reaction indicator value: **6.5**
Nutrient indicator value: **4.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.88**
Disturbance frequency (herb layer): **1.49**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.66**
Grazing pressure: **0.19**
Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Forest, Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz