

Agrostis canina

Habitus and growth type

Plant height [m]: **0.37**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **28.65**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5**

Moisture indicator value: **8.4**

Reaction indicator value: **3.1**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.05**

Disturbance frequency (herb layer): **1.56**

Disturbance severity: **0.4**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.44**

Grazing pressure: **0.22**

Soil disturbance: **0.17**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q22 Poor fen](#), [Q23 Relict mire of Mediterranean mountains](#)

Constant species of EUNIS habitats: [Q22 Poor fen](#), [Q23 Relict mire of Mediterranean mountains](#), [Q24 Intermediate fen and soft-water spring mire](#), [Q42 Extremely rich moss-sedge fen](#), [R34 Submediterranean moist meadow](#), [R35 Moist or wet mesotrophic to eutrophic hay meadow](#), [R37 Temperate and boreal moist or wet oligotrophic grassland](#), [T16 Broadleaved mire forest on acid peat](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **6**