

Achillea clypeolata

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.11**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **7.1**

Moisture indicator value: **3.2**

Reaction indicator value: **7.2**

Nutrient indicator value: **2.7**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.97**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.25**

Grazing pressure: **0.33**

Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N1E Black Sea broad-leaved coastal dune](#)

[forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**