Achillea erba-rotta

Habitus and growth type

Plant height [m]: **0.2**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **22.22**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.34**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **1.3**
Moisture indicator value: **4.8**
Reaction indicator value: **2.5**
Nutrient indicator value: **1.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.15**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.52**
Disturbance severity: **0.36**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.02**
Grazing pressure: **0.22**
Soil disturbance: **0.14**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **U22 Temperate high-mountain siliceous scree**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz