

# *Achillea maritima*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Chamaephyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.89**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **9**

Moisture indicator value: **3.7**

Reaction indicator value: **6.5**

Salinity indicator value: **3.8**

### Disturbance indicator values

Disturbance frequency: **2.04**

Disturbance frequency (herb layer): **2.07**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.41**

Mowing frequency: **0**

Grazing pressure: **0.12**

Soil disturbance: **0.73**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [JD \(AMM\) \*Ammophiletea arundinaceae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [N14 Mediterranean, Macaronesian and Black Sea shifting coastal dune](#), [N22 Mediterranean and Black Sea coastal shingle beach](#)

Constant species of EUNIS habitats: [N14 Mediterranean, Macaronesian and Black Sea shifting coastal dune](#), [N22 Mediterranean and Black Sea coastal shingle beach](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle**