

Achillea micrantha

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.8**

Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **1.67**

Disturbance frequency (herb layer): **1.79**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.41**

Mowing frequency: **0.35**

Grazing pressure: **0.22**

Soil disturbance: **0.39**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**BD \(PYR\) *Pyrolo-Pinetea sylvestris*,
CJ \(COR\) *Koelerio-Corynephorotea canescentis***](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R11 Pannonian and Pontic sandy steppe**](#)

Constant species of EUNIS habitats: [**R11 Pannonian and Pontic sandy steppe**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**