Achillea pastricensis

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.9**
Temperature indicator value: **7.1**
Moisture indicator value: **3.4**
Reaction indicator value: **6.7**
Nutrient indicator value: **2.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.96**
Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.38**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.03**
Grazing pressure: **0.31**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FR (DAP) *Daphno-Festucetea***, **KC (ASP) *Asplenietea trichomanis***

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl.**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

rock and scree)

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz