

Achillea pastricensis

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.9**

Temperature indicator value: **7.1**

Moisture indicator value: **3.4**

Reaction indicator value: **6.7**

Nutrient indicator value: **2.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.96**

Disturbance frequency (herb layer): **1.97**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.03**

Grazing pressure: **0.31**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FR \(DAP\) *Daphno-Festucetea*](#), [KC \(ASP\) *Asplenietea trichomanis*](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl.**

rock and scree)