Aira elegantissima

Habitus and growth type

Plant height [m]: **0.21**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.05**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**
Moisture indicator value: **2.5**
Nutrient indicator value: **1.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.84**
Disturbance frequency (herb layer): **1.77**
Disturbance severity: **0.56**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.35**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea*, FI (TUB) *Helianthemetea guttati***

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz