

# *Ajuga chamaepitys*

## Habitus and growth type

Plant height [m]: **0.11**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Spinescence: **not spinescent**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **20.28**

## Flower

Flowering period: **March-October**

## Fruit, seed and dispersal

Seed mass [mg]: **1.54**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.5**

Moisture indicator value: **3.6**

Reaction indicator value: **8.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.47**

Disturbance frequency (herb layer): **1.98**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.63**

Mowing frequency: **0.59**

Grazing pressure: **0.3**

Soil disturbance: **0.42**

## **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality: 7

Continentality amplitude: 7