

# *Alisma lanceolatum*

## Habitus and growth type

Plant height [m]: **0.43**

Life span: **Perennial**

Life form: **Hemicryptophyte, Hydrophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **28.33**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.43**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **10.1**

Reaction indicator value: **6.6**

Nutrient indicator value: **5.9**

Salinity indicator value: **0.1**

### Disturbance indicator values

Disturbance frequency: **1.42**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.2**

Grazing pressure: **0.13**

Soil disturbance: **0.14**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*OD \(PHR\) \*Phragmito-Magnocaricetea\*\*\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*V14 Inundated or inundatable cropland, including rice fields\*\*](#)

Constant species of EUNIS habitats: [\*\*V14 Inundated or inundatable cropland, including rice fields\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

## Distribution

Continentality: **7**

Continentality amplitude: **8**