

Alliaria petiolata

Habitus and growth type

Plant height [m]: **0.5**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **40.83**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.58**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.2**

Temperature indicator value: **5.9**

Moisture indicator value: **5.4**

Reaction indicator value: **7.3**

Nutrient indicator value: **8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.15**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.07**

Grazing pressure: **0.19**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#),
[HA \(POP\) *Alno glutinosae-Populetea albae*](#), [QG \(EPI\) *Epilobietea angustifolii*](#)

EUNIS habitat

Constant species of EUNIS habitats: [T11 Temperate *Salix* and *Populus* riparian forest](#), [T13 Temperate hardwood riparian forest](#), [T1H Broadleaved deciduous plantation of non site-native trees](#), [T38 Mediterranean montane *Cedrus* forest](#),
[V39 Mesic perennial anthropogenic herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Synanthropic**

Distribution

Continentality amplitude: **6**