

Allium cupani

Habitus and growth type

Plant height [m]: **0.17**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.32**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **4**

Moisture indicator value: **3**

Reaction indicator value: **5**

Nutrient indicator value: **5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.15**

Disturbance frequency (herb layer): **2.15**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.15**

Grazing pressure: **0.37**

Soil disturbance: **0.18**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline),
Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**