

Allium ericetorum

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.62**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **5.5**

Moisture indicator value: **5**

Reaction indicator value: **5**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.47**

Disturbance frequency (herb layer): **1.38**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.2**

Grazing pressure: **0.26**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) *Calluno-Ulicetea*](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Heathland, Sparsely vegetated (incl. rock and scree)**