

Allium flavum

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.35**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **6.5**

Moisture indicator value: **3**

Reaction indicator value: **6.8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.88**

Disturbance frequency (herb layer): **1.65**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.22**

Grazing pressure: **0.32**

Soil disturbance: **0.18**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R17 Heavy-metal dry grassland of the Balkans**](#)

Constant species of EUNIS habitats: [**R16 Perennial rocky grassland of Central and South-Eastern Europe**](#), [**R17 Heavy-metal dry grassland of the Balkans**](#), [**S36 Low steppic scrub**](#), [**U3A Temperate ultramafic inland cliff**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**,
Oromediterranean scrub, **Scrub**, **Forest**, **Sparingly vegetated (incl. rock and scree)**