

# *Allium pallens*

## Habitus and growth type

Plant height [m]: **0.53**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **1.62**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **9.3**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.53**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.35**

Grazing pressure: **0.38**

Soil disturbance: **0.19**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [FF \(LYG\) \*Lygeo sparti-Stipetea tenacissimae\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**