

Allium roseum

Habitus and growth type

Plant height [m]: **0.39**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.83**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **8.6**

Moisture indicator value: **3.4**

Reaction indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.78**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.44**

Grazing pressure: **0.28**

Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**DA \(FES\) Festuco-Brometea**](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**