# *Allium triquetrum*

### Habitus and growth type

Plant height [m]: **0.25**
Life span: **Perennial**
Life form: **Geophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **23.9**

### Flower

Flowering period: **January-April, November-December**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **6**
Temperature indicator value: **8.5**
Moisture indicator value: **4**
Reaction indicator value: **6.7**
Nutrient indicator value: **6.7**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.22**
Disturbance frequency (herb layer): **1.33**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0.08**
Grazing pressure: **0.24**
Soil disturbance: **0.14**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea*, QG (EPI) *Epilobietea angustifolii***

### Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*