

Allium ursinum

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **37.25**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.04**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **7.4**

Nutrient indicator value: **7.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.16**

Mowing frequency: **0.01**

Grazing pressure: **0.18**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#),
[HA \(POP\) *Alno glutinosae-Populetea albae*](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

Distribution

Continentality: **4**

Continentality amplitude: **4**