

# *Amelanchier lamarckii*

## Habitus and growth type

Plant height [m]: **9.25**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **11.34**

## Fruit, seed and dispersal

Seed mass [mg]: **3.89**

Dispersal mode: **Endozoochory, Anthropochory**

Dispersal distance class: **6, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.7**

Temperature indicator value: **6.8**

Moisture indicator value: **4.7**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.08**

Disturbance frequency (herb layer): **1.27**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.16**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.2**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CD \(RHA\) Crataego-Prunetea](#)

Broad habitat

Occurrence in broad habitats: **Scrub**

## **Distribution**

Continentality: **5**

Continentality amplitude: **4**