

Anagyris foetida

Habitus and growth type

Plant height [m]: **3.25**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Specific leaf area [mm²/mg]: **10.48**

Flower

Flowering period: **January-April**

Fruit, seed and dispersal

Seed mass [mg]: **358.9**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **9**

Moisture indicator value: **3.3**

Reaction indicator value: **7**

Nutrient indicator value: **5.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.27**

Disturbance frequency (herb layer): **1.47**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.04**

Grazing pressure: **0.28**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FA \(QUI\) Quercetea ilicis**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**