

# *Androsace septentrionalis*

## Habitus and growth type

Plant height [m]: **0.11**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.53**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.24**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **5.7**

Moisture indicator value: **3**

Reaction indicator value: **5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.6**

Disturbance frequency (herb layer): **2.07**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.53**

Mowing frequency: **0.67**

Grazing pressure: **0.32**

Soil disturbance: **0.25**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Coryneporetea canescentis\*](#), [CK \(SED\) \*Sedo-Scleranthetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [U35 Boreal and Arctic base-rich inland cliff](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **7**

Continentality amplitude: **6**