

Anemone sylvestris

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm²/mg]: **18.32**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.56**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.6**

Moisture indicator value: **3.5**

Reaction indicator value: **7.4**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.7**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.47**

Grazing pressure: **0.28**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#), [CL \(GER\) Trifolio-Geranietea sanguinei](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: 7

Continentality amplitude: 5