

Anisantha madritensis

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **32**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **2.56**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **2.8**

Reaction indicator value: **7.3**

Nutrient indicator value: **6**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.04**

Disturbance frequency (herb layer): **1.8**

Disturbance severity: **0.79**

Disturbance severity (herb layer): **0.72**

Mowing frequency: **0.5**

Grazing pressure: **0.25**

Soil disturbance: **0.4**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) *Chenopodietea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V32 Mediterranean subnitrophilous annual grassland](#)

Constant species of EUNIS habitats: [V32 Mediterranean subnitrophilous annual grassland](#), [V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**