Anisantha sterilis

Habitus and growth type

Plant height [m]: **0.39**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **31.82**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **8.24**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **6.3**
Moisture indicator value: **3.9**
Reaction indicator value: **6.7**
Nutrient indicator value: **6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.78**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.65**
Disturbance severity: **0.73**
Disturbance severity (herb layer): **0.61**
Mowing frequency: **0.46**
Grazing pressure: **0.2**
Soil disturbance: **0.42**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea***

EUNIS habitat

Constant species of EUNIS habitats: **T3D Mediterranean Cupressaceae forest**, **V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation**, **V37 Annual anthropogenic herbaceous vegetation**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz