Antinoria agrostidea

Habitus and growth type

Life span: **Annual**
Life form: **Hydrophyte, Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6**
Moisture indicator value: **9**
Reaction indicator value: **5**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.37**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.12**
Disturbance severity (herb layer): **0.12**
Mowing frequency: **0.02**
Grazing pressure: **0.01**
Soil disturbance: **0.02**

Habitat and sociology

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Aquatic, Wetland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz