# *Antirrhinum latifolium*

### Habitus and growth type

Plant height [m]: **0.75**
Life span: **Perennial**
Life form: **Hemicryptophyte**

### Flower

Flowering period: **April-September**

### Fruit, seed and dispersal

Seed mass [mg]: **0.15**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **8**
Moisture indicator value: **2**
Nutrient indicator value: **1**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.35**
Disturbance frequency (herb layer): **1.38**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.05**
Grazing pressure: **0.32**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **KC (ASP) *Asplenietea trichomanis***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*