

# *Antirrhinum majus*

## Habitus and growth type

Plant height [m]: **0.58**

Life span: **Annual, Perennial**

Life form: **Chamaephyte, Hemicryptophyte, Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **17.2**

## Flower

Flowering period: **January-November**

## Fruit, seed and dispersal

Seed mass [mg]: **0.11**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.1**

Moisture indicator value: **3.8**

Reaction indicator value: **7.1**

Nutrient indicator value: **5.8**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **0.61**

Disturbance frequency (herb layer): **1.05**

Disturbance severity: **0.6**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.25**

Grazing pressure: **0.21**

Soil disturbance: **0.23**

## **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality: **2**

Continentality amplitude: **2**