Aquilegia dichroa

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **9**
Moisture indicator value: **11**
Reaction indicator value: **5**
Nutrient indicator value: **5**

Disturbance indicator values

Disturbance frequency: **0.3**
Disturbance frequency (herb layer): **1.47**
Disturbance severity: **0.62**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.1**
Grazing pressure: **0.2**
Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz