

Arabis auriculata

Habitus and growth type

Plant height [m]: **0.22**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.05**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **7**

Moisture indicator value: **3.4**

Reaction indicator value: **7.4**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.07**

Disturbance frequency (herb layer): **1.8**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.19**

Grazing pressure: **0.34**

Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*](#), [FI \(TUB\) *Helianthemetea guttati*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **7**

Continentality amplitude: **5**